$See \ discussions, stats, and author \ profiles \ for \ this \ publication \ at: \ https://www.researchgate.net/publication/260367809$

MIXED DENTITION SPACE ANALYSIS: A REVIEW

Article · December 2012

CITATIONS		READS	
6		25,371	
4			
4 autors:			
	Arsalan Wahid	Q	Sidra Butt
	The University of Faisalabad		COMSATS University Islamabad
	9 PUBLICATIONS 137 CITATIONS		6 PUBLICATIONS 94 CITATIONS
	SEE PROFILE		SEE PROFILE
Q	Saima Chaudhry	Q	Afifa Ehsan
	University of Health Sciences Lahore		Sheikh Zayed Medical College & Hospital
	18 PUBLICATIONS 456 CITATIONS		18 PUBLICATIONS 128 CITATIONS
	SEE PROFILE		SEE PROFILE

MIXED DENTITION SPACE ANALYSIS: A REVIEW

¹SIDRA BUTT, M.Phil SCHOLAR ²SAIMA CHAUDHRY, PhD (ORAL PATHOLOGY) ³MARIYAH JAVED, M.Phil ⁴ARSALAN WAHID, M.Phil SCHOLAR ⁵AFIFA EHSAN, M.Phil SCHOLAR ⁶SOBIA MALIK M.Phil ⁷AYYAZ ALI KHAN, PhD

ABSTRACT

During mixed dentition period discrepancy between spaces available and required in the arch is predicted so that teeth can erupt freely with good alignment. Mixed dentition space analysis is done for this purpose in which size of unerupted canine and premolars is predicted. Accurate prediction is very important regarding correct orthodontic diagnosis and treatment planning. Prediction equation and or tables based on size of already erupted permanent teeth are the most widely used methods. Tooth size varies in different racial and ethnic groups and among genders so prediction equation and tables specific to each population for both male and female must be made. Mandibular incisors; combination of mandibular incisors and mandibular 1st molars; and mandibular incisors and maxillary 1st molars have been used worldwide. Recent studies have reported that mandibular incisors are not the best predictors and sum of mandibular incisors and maxillary 1st molars are even more accurate predictor for the estimation of size of unerupted teeth. In Pakistani population combination of mandibular incisors and maxillary 1st molar have not been employed so study must be conducted to estimate which combination of group of teeth is more accurate for the estimation of size of unerupted canine and premolars in local population.

Keywords: Mixed dentition, space analysis, mesiodistal width, unnerupted, prediction equation.

INTRODUCTION AND LITERATURE REVIEW

Dental occlusion is the way in which maxillary and mandibular teeth come into contact with each other. It can be static when the jaw is in rest position or centric in which maxillary and mandibular teeth are in maximum intercuspation position.¹ For all orthodontic patients achievement ideal dental occlusion is the main therapeutic goal.²

DENTAL MALOCCLUSION

Malocclusion is any deviation from normal occlusion. Dental malocclusion can occur as a result of genetic factors, environmental factors or combination of both factors. Most of them are the result of an imbalance between teeth size and arch sizes with which teeth are associated.³ Dental malocclusion arise during mixed dentition period and if they are managed timely they can be reduced in severity or even removed entirely.⁴

Corresponding Author ¹M.Phil. Scholar, Department of Oral Health Sciences, Shaikh Zayed Medical Complex, Lahore. Mobile no: 03454653213, E-mail: sid_nmc@hotmail.com

² Lecturer, Cell: 03334416740, E-mail: chaudhry_s@iadsr.edu.pk

³ M.Phil. Scholar, Mobile no: 03314090878, E-mail: mariyah.javed@hotmail.com

⁴ M.Phil. Scholar, Mobile no: 03216685228, E-mail: dr.arsalanmalik@gmail.com

⁵ M.Phil. Scholar, Mobile no: 03004668379, E-mail: afifaehsan@gmail.com

⁶ Assistant Professor

⁷ Prof and Head of Department of Oral Health Sciences, Shaikh Zayed Medical Complex, Lahore. Mobile No: 03008454981,

Mixed Dentition Period and Its Importance in Interceptive Orthodontics

Mixed dentition period is between six to twelve years of age during which both deciduous and permanent teeth are present. During this period one can make accurate and specific prediction of future dental development and also can assess whether there will be spacing or crowding of teeth in the dental arch.⁵ In interceptive orthodontics treatment planning, it is important to predict space required and available for unerupted canine and premolars in the arch and this determination of tooth size must be done before eruption of canine and premolars by a method called Mixed Dentition Space Analysis (MDSA).⁶

Mixed Dentition Space Analysis

In mixed dentition space analysis, mesiodistal width of unerupted canine and premolars is predicted so that discrepancy between space available and space required for these teeth in the dental arch can be determined.⁷⁻¹² This also helps to determine whether sufficient space is available for posterior teeth in the arch so that they can erupt freely with good alignment.¹³

This analysis is very important regarding orthodontic diagnosis and treatment planning.¹⁴ It helps in determining whether treatment plan is going to involve serial extractions, eruption guidance, regaining of space, space maintenance, or simple observation of patient for the time period is required.^{15,16} Accurate prediction of size of unerupted posterior teeth is important in order to establish a good orthodontic treatment plan.^{17,18}

METHODOLOGY

A review of literature reveals three main approaches used to estimate mesiodistal width of unerupted canine and premolars in mixed dentition patients: 1) Direct measurement of the teeth from dental radiograph (periapical and cephalometric)^{8,9}; 2) Using prediction equations and tables based on measurements of deciduous teeth⁸ or other erupted permanent teeth^{7, 12}; 3) Combination of both methods.^{8,19-21}

i. Direct Measurement from Radiograph

Width of permanent canine and premolars (first and second) is measured directly from dental radio-

graphs (periapical and cephalometric). Several studies have reported methods to improve accuracy for the estimation of size of unerupted teeth by using dental radiographs.²¹⁻²⁶

ii. Use of Prediction Equation and/ or Table

Prediction equation and or tables based on of mesiodistal width of already erupted permanent teeth are used to estimate size of unerupted canine and premolars. The most commonly used are Moyer's probability tables and Tanaka and Johnston's regression equations.^{7,12}

iii. Combination of Radiograph and Prediction Equation

In combined method, mesiodistal width of permanent canine and premolars is predicted from sum width of first and second premolars measured from radiograph plus width of erupted mandibular central and lateral incisor of the same quadrant.^{21,27} It was first recommended by Hixon and Old Father (1958). He established regression equation to predict size of unerupted mandibular canine and premolars from radiographic measurement of mandibular premolars.^{21,28} Oldfather's method was modified by Stahle (1959) by including the sizes of the mandibular permanent incisor to predict mesiodistal width of canine and premolars.²⁷ Staley, R. N. and P. E. Kerber (1980) revised Hixon and Old Father and developed prediction equation and made a graph for the estimation of mesiodistal width of unerupted caine and premolars.²⁰ Several researchers predicted mesiodistal width of unerupted canine and premolars in Caucasian population by a combination of measurements of teeth from dental radiograph and regression analysis done on dental casts and they found it to be more accurate in their respective population.^{29, 30}

Radiographic Vs Prediction Equation and / or Table

Accurate radiographic prediction requires good quality radiographic technique and equipment. Radiograph gives two dimensional image of object so it may give false measurement when teeth may be rotated in their bony crypt.¹⁶ It also have disadvantage of exposure to increase radiation dose. So prediction equation and or tables based on mesiodistal width of already erupted permanent teeth are the most widely used methods. The most commonly used are Moyer's probability tables and Tanaka and Johnston's regression equations. 7,12

Moyers Probability Method

Moyers (1958) developed two probability tables to predict mesiodistal width of unerupted canine & premolars during mixed dentition based on sum of mesiodistal width of four mandibular permanent incisors.¹² This is the most widely used for mixed dentition because of the following advantages: 1) Minimal known systematic error. 2) Can be used by beginner and expert with equal reliability. 3) Less time consuming. 4) doesn't require specific equipment or exposure to radiographs. 5) Applicable to both upper and lower dental arches. 5) Best done on dental cast bit it can be done in mouth with reasonable accuracy. Moyers method is applied most frequently by many populations.^{7,12,23,31,32} Profit and Ackerman also found Moyers method to be satisfactory.³³

Tanaka and Johnston Method

Tanaka and Johnston (1974) developed probability equation to predict mesiodistal width of unerupted canine & premolars by using sum of width of four mandibular incisors.⁷ This is simple, easy, relatively noninvasive and accurate, applicable for both dental arches and genders. This method has also been used widely with an acceptable accuracy for both dental arches and in both genders.^{7,34-36}

Racial and Ethnic Difference in Tooth Size

Literature shows that tooth size varies among different racial and ethnic groups. This variation in tooth size is related to genetic, epigenetic and environmental factors.^{37,38}

Since Moyer and Tanaka and Johnston prediction equation and tables were developed for North American Caucasian children. So its applicability in population of other racial and ethnic origins has been studied and doubted by many researchers.^{4,6,14,18,36,39-62}

Gender Difference in Tooth Size

These methods were not developed for both genders separately while literature shows that tooth size varies among both males and females with males having larger teeth than females.^{6,14,43,45,46,54,59,60,63} Many authors reported that sex factor should also be included as an additional predictor for the estimation of MDW sum of unerupted canine and premolars based on sexual dimorphism on teeth size. So regression equation and or tables must be made separate for both males and females.^{11,45,63}

Combination of Group of Teeth Used as Predictor

Different combinations of teeth have been used as predictor for the estimation of size of unerupted canine and premolars in mixed dentition space analysis.

• Sum of Four Mandibular Incisors

Sum of mesiodistal width of four mandibular incisors has been most widely used to develop regression equation to accurately predict size of unerupted canine & premolars specific to their own population by many researchers.^{16,39,41,44,46-51,53,54,57,59,61,64}

• Other Combinations of Group of Teeth Used as Predictor

In addition to mandibular incisors, combinations of other erupted permanent teeth (e.g mandibular first molar, maxillary first molar and maxillary incisors) also have been used by some researchers to accurately predict size of unerupted canine & premolars. Fonseca (1961) firstly introduced an additional combination of teeth (sum of maxillary first molar and maxillary four permanent incisors) to develop a multiple regression equation.⁶⁵ Recent studies have reported that sum of mesiodistal width of permanent mandibular incisors is not the best predictor for estimating mesiodistal width of unerupted permanent canine & premolars.^{11,63,66,67} so other teeth along with incisors also have been evaluated to predict mesiodistal width of unerupted canine & premolars.

- Sum of mesiodistal width of permanent mandibular incisors and mandibular first molar has been used as predictor in Brazilian population.⁶⁷
- Sum of mesiodistal width of maxillary central incisor and mandibular first molar is found to be the best predictor to estimate mesiodistal width of permanent canine & premolars in Spanish and Egyptian population.^{52,58}

- Bernabe E, Flores Mir-C (2005) reported that sum of maxillary and mandibular central incisor and maxillary 1st molar is more accurate predictor for estimating mesiodistal width of unerupted canine & premolars in Peru population.¹¹
- Sum of mesiodistal width of maxillary first molar and mandibular incisors is found to be the best predictor to estimate mesiodistal width of unerupted canine & premolars by most of the researchers in Syrian, Croatia and Italian population.^{63,66,68}

Nourallah (2001) developed a more accurate prediction equation and table in Syrian population based on sum of mesiodistal width of maxillary first molar and mandibular central incisors to predict size of unerupted canine & premolar.⁶⁶ Legovic et al (2003) developed regression equation based on mesiodistal and buccolingual width of permanent incisors and maxillary first molar to estimate size of unerupted canine and premolars in a population sample of Croatia.⁶³ Cattaneo C et al (2010) studied combination of group of teeth to predict size of unerupted canine and premolars with most common groups of teeth including four mandibular incisors; mandibular incisors and mandibular first molar; mandibular incisors and maxillary first molar and concluded that sum of four mandibular incisors and maxillary 1st molar is the best predictor for estimating mesiodistal width of unerupted permanent canine & premolars in Italian sample.68

Studies Conducted in Pakistan

Applicability of Moyers and Tanaka method has been tested by some researchers and they found that these methods do not accurately predict size of unerupted canine and premolars in our population.^{4,59} Rasool G et al (2008) checked applicability of regression equation developed by Melgaco to estimate mesiodistal width of unerupted mandibular canine and premolars from sum of mandibular incisors and first molars. They concluded that this equation is not suitable in males while it can be applied to females in Islamabad population (100 males, 100females).⁶⁹ In 2011, prediction equation to estimate MDW of unerupted canine and premolars was developed from sum of MDW mandibular incisors and mandibular first molars separate for both males and females in the mandibular arch in Karachi population (106 males, 182 females). 69

Bherwani et al (2011) developed regression equation to predict mesiodistal width of maxillary and mandibular canine and premolars from sum of mesiodistal width of mandibular incisors with significant correlation for maxillary arch (r=0.65) and mandibular arch (r=0.59) on Karachi population (100 boys, 100 girls).¹⁶ They reported no significant gender difference in tooth sizes while other researchers have reported larger tooth size in males as compared to females in the country.^{4,59,69,70} In this regard prediction equation separate for both males and females and both arches must be developed to enhance accuracy and treatment planning in mixed dentition space analysis in Pakistani population.

Accurate prediction of mesiodistal width of unerupted permanent teeth is required to determine tooth size and arch length discrepancies in mixed dentition. Inaccurate mixed dentition space analysis may lead to wrong extraction decision and poor patient's facial profile.⁵⁴ As patient's demand to seek an early orthodontic treatment is increasing therefore it is important for orthodontists to predict any discrepancy in arch space in advance and initiate appropriate treatment in a timely manner.

CONCLUSION

Mandibular incisors alone; and combination of Mandibular incisors and mandibular first molar, have been used to estimate the mesiodistal width of unerupted canine & premolars in Pakistani population, but combination of maxillary first molar and mandibular incisors, which has been found to be better than the above mentioned group of teeth for estimation of mesiodistal width of canine & premolar, has not been validated on Pakistani population. There is a need to conduct large scale study on a local population sample to evaluate which combination of group of teeth gives more accurate prediction for the estimation of size of unerupted canine and premolars so that discrepancy between tooth size and arch perimeter can be diagnosed early and appropriate treatment can be initiated in time. This will prevent patient from developing adult malocclusion leading to prolonged and expensive future dental management.

REFERENCES

- 1 M.S. Rani, editor. Textbook of orthodontics. 3 ed. Chennai, India: Rajender K. Arya for.; 2001.
- 2 Oltramari PV, Conti AC, Navarro Rde L, Almeida MR, Almeida-Pedrin RR, Ferreira FP. Importance of occlusion aspects in the completion of orthodontic treatment. Braz Dent J. 2007; 18(1): 78-82.
- 3 Lima Martinelli F, Martinelli de Lima E, Rocha R, Souza Tirre-Araujo M. Prediction of lower permanent canine and premolars width by correlation methods. Angle Orthod. 2005 Sep; 75(5): 805-08.
- 4 Mengal N, Afzal A. Mixed dentition analysis for Pakistani population. Journal of Surgery Pakistan International. 2004; 9(1): 10-14.
- 5 Hunter WS. Application of analysis of crowding and spacing of the teeth. Dent Clin North Am. 1978 Oct; 22(4): 563-77.
- 6 Ahluwalia P, Jodhka S, Thomas AM. Prediction of Mesio-distal width of Canines and premolars in a sample of north Indian population. Indian Journal of Dental Advancements. 2011; 3.
- 7 Tanaka MM, Johnston LE. The prediction of the size of unerupted canines and premolars in a contemporary orthodontic population. J Am Dent Assoc. 1974 Apr; 88(4): 798-801.
- 8 Bishara SE, Staley RN. Mixed-dentition mandibular arch length analysis: a step-by-step approach using the revised Hixon-Oldfather prediction method. Am J Orthod. 1984 Aug; 86(2): 130-35.
- 9 de Paula S, Almeida MA, Lee PC. Prediction of mesiodistal diameter of unerupted lower canines and premolars using 45 degrees cephalometric radiography. Am J Orthod Dentofacial Orthop. 1995 Mar; 107(3): 309-14.
- 10 Schirmer UR, Wiltshire WA. Orthodontic probability tables for black patients of African descent: mixed dentition analysis. Am J Orthod Dentofacial Orthop. 1997 Nov; 112(5): 545-51.
- 11 Bernabe E, Flores-Mir C. Are the lower incisors the best predictors for the unerupted canine and premolars sums? an analysis of a Peruvian sample. Angle Orthod. 2005 Mar; 75(2): 202-07.
- 12 Moyer, editor. handbook of orthodontics. 4th ed. chicago: year book medical publisher; 1958: p. 577.
- 13 Kirschen RH, O'Higgins EA, Lee RT. The Royal London Space Planning: an integration of space analysis and treatment planning: Part II: The effect of other treatment procedures on space. Am J Orthod Dentofacial Orthop. 2000 Oct; 118(4): 456-61.
- 14 Al-Bitar ZB, Al-Omari IK, Sonbol HN, Al-Ahmad HT, Hamdan AM. Mixed dentition analysis in a Jordanian population. Angle Orthod. 2008 Jul; 78(4): 670-75.
- 15 Smith HP, King DL, Valencia R. A comparison of three methods of mixed-dentition analyses. J Pedod. 1979 Summer; 3(4): 291-302.
- 16 Bherwani AK, Fida M. Development of a prediction equation for the mixed dentition in a Pakistani sample. Am J Orthod Dentofacial Orthop. 2011 Nov; 140(5): 626-32.
- 17 Bolton W. The clinical application of a tooth Size analysis. AmJ Orthod. 1962; 48: 504-29.
- 18 Bolton W. Disharmony in tooth size and its relation to the analysis in treatment of malocclusion. Angle Orthod. 1958; 28: 113-30.

- 19 Staley RN, O'Gorman TW, Hoag JF, Shelly TH. Prediction of the widths of unerupted canines and premolars. J Am Dent Assoc. 1984 Feb; 108(2): 185-90.
- 20 Staley RN, Kerber PE. A revision of the Hixon and Oldfather mixed-dentition prediction method. Am J Orthod. 1980 Sep; 78(3): 296-302.
- 21 Hixon EH OR. Estimation of the sizes of unerupted cuspid and bicuspid teeth. Angle Orthod. 1958; 28: 236-40.
- 22 Nance HN. The limitations of orthodontic treatment . Mixed dentition: diagnosis and treatment. Am J Orthod Oral Surg. 1947 Apr; 33(4): 177-223.
- 23 Ballard ML, Wylie WL. Mixed dentition case analysis, estimating size of unerupted permanent teeth. Am J Orthod. 1947 Nov; 33(11): 754-59.
- 24 Cohen M. Recognition of the developing Malocclusion. Dental Clinics of North America. 1959; 6: 299–311.
- 25 Foster RR WW. Arch length deficiency in the mixed dentition. American Journal of Orthodontics and Dentofacial Orthopedics. 1958; 44: 464–76.
- 26 Moorrees CF, Reed RB. Correlations among Crown Diameters of Human Teeth. Arch Oral Biol. 1964 Nov-Dec; 9: 685-97.
- 27 Stahle. Determination of mesiodistal crown width of the unerupted permanent cuspids and bicuspids. Helv Orthod Acta. 1959; 3: 14-9.
- 28 RH. O. Estimation of the sum of the widths of unerupted mandible cuspid, first bicuspid, and second bicuspid. Am J Orthod. 1957(43): 788-89.
- 29 Staley RN, Hoag JF. Prediction of the mesiodistal widths of maxillary permanent canines and premolars. Am J Orthod. 1978 Feb; 73(2): 169-77.
- 30 Staley RN, Shelly TH, Martin JF. Prediction of lower canine and premolar widths in the mixed dentition. Am J Orthod. 1979 Sep; 76(3): 300-09.
- 31 Bishara SE, Hoppens BJ, Jakobsen JR, Kohout FJ. Changes in the molar relationship between the deciduous and permanent dentitions: a longitudinal study. Am J Orthod Dentofacial Orthop. 1988 Jan; 93(1): 19-28.
- 32 Jensen E, Kai-Jen Yen P, Moorrees CF, Thomsen SO. Mesiodistal crown diameters of the deciduous and permanent teeth in individuals. J Dent Res. 1957 Feb; 36(1): 39-47.
- 33 Proffit W AJ. Contemporary Orthodontics. St Louis, MO1993.
- 34 Foster HR. An analysis of the developing dentition. J Am Dent Assoc. 1951 Apr; 42(4): 376-87.
- 35 HR F. Arch length deficiency in the mixed dentition. Am JOrthod. 1958; 44: 464-74.
- 36 Gardner RB. A comparison of four methods of predicting arch length. Am J Orthod. 1979 Apr; 75(4): 387-98.
- Townsend G, Bockmann M, Hughes T, Brook A. Genetic, environmental and epigenetic influences on variation in human tooth number, size and shape. Odontology. 2012 Jan; 100(1): 1-9.
- 38 Brook AH, Griffin RC, Townsend G, Levisianos Y, Russell J, Smith RN. Variability and patterning in permanent tooth size of four human ethnic groups. Arch Oral Biol. 2009 Dec; 54 Suppl 1: S79-85.

- 39 Abu Alhaija ES, Qudeimat MA. Mixed dentition space analysis in a Jordanian population: comparison of two methods. Int J Paediatr Dent. 2006 Mar; 16(2): 104-10.
- 40 Alessandri Bonetti G, Verganti S, Zanarini M, Bonetti S, Gatto MR. Mixed dentition space analysis for a northern Italian population: new regression equations for unerupted teeth. Prog Orthod. 2011 Nov; 12(2): 94-99.
- 41 al-Khadra BH. Prediction of the size of unerupted canines and premolars in a Saudi Arab population. Am J Orthod Dentofacial Orthop. 1993 Oct; 104(4): 369-72.
- 42 Altherr ER, Koroluk LD, Phillips C. Influence of sex and ethnic tooth-size differences on mixed-dentition space analysis. American Journal of Orthodontics and Dentofacial Orthopedics. 2007; 132(3): 332-39.
- 43 Arslan SG, Dildes N, Kama JD, Genc C. Mixed-dentition analysis in a Turkish population. World J Orthod. 2009 Summer; 10(2):135-40.
- 44 Diagne F, Diop-Ba K, Ngom PI, El Boury O. Mixed dentition analysis in a Moroccan population. Odontostomatol Trop. 2004 Dec; 27(108): 5-10.
- 45 Hashim HA, Al-Shalan TA. Prediction of the size of un-erupted permanent cuspids and bicuspids in a Saudi sample: a pilot study. J Contemp Dent Pract. 2003 Nov 15; 4(4): 40-53.
- 46 Jaroontham J, Godfrey K. Mixed dentition space analysis in a Thai population. Eur J Orthod. 2000 Apr; 22(2): 127-34.
- 47 Khan MI, Seedat AK, Hlongwa P. Tooth width predictions in a sample of Black South Africans. SADJ. 2007 Jul; 62(6): 244, 246-49.
- 48 nik Tahere H, Majid S, Fateme M, Kharazi f, Javad M. Predicting the size of unerupted canines and premolars of the maxillary and mandibular quadrants in an Iranian population. J Clin Pediatr Dent. 2007 Fall; 32(1): 43-47.
- 49 Ling JY, Wong RW. Tanaka-Johnston mixed dentition analysis for southern Chinese in Hong Kong. Angle Orthod. 2006 Jul; 76(4): 632-36.
- 50 Philip NI, Prabhakar M, Arora D, Chopra S. Applicability of the Moyers mixed dentition probability tables and new prediction aids for a contemporary population in India. Am J Orthod Dentofacial Orthop. 2010 Sep; 138(3): 339-45.
- 51 Peng H, Wang X, Chen K. [The predication equation of the permanent canine and premolar crown]. Hua Xi Kou Qiang Yi Xue Za Zhi. 2000 Feb; 18(1): 55-57.
- 52 Paredes V, Gandia JL, Cibrian R. A new, accurate and fast digital method to predict unerupted tooth size. Angle Orthod. 2006 Jan; 76(1): 14-9.
- 53 Diagne F, Diop-Ba K, Ngom PI, Mbow K. Mixed dentition analysis in a Senegalese population: elaboration of prediction tables. Am J Orthod Dentofacial Orthop. 2003 Aug; 124(2): 178-83.
- 54 Durgekar SG, Naik V. Evaluation of Moyers mixed dentition analysis in school children. Indian J Dent Res. 2009 Jan-Mar; 20(1): 26-30.

- 55 Ferguson FS, Macko DJ, Sonnenberg EM, Shakun ML. The use of regression constants in estimating tooth size in a Negro population. Am J Orthod. 1978 Jan; 73(1): 68-72.
- 56 Frankel HH, Benz EM. Mixed dentition analysis for black Americans. Pediatr Dent. 1986 Sep; 8(3): 226-30.
- 57 Jaiswal AK, Paudel KR, Shrestha SL, Jaiswal S. Prediction of space available for unerupted permanent canine and premolars in a Nepalese population. J Orthod. 2009 Dec; 36(4): 253-59.
- 58 Ibrahim IA, Elkateb MA, Wahba NA, El Harouny NM. A digital method to predict the mesiodistal widths of canines and premolars in an Egyptian sample. J Clin Pediatr Dent. 2011 Summer; 35(4): 421-27.
- 59 Memon S, Fida M. Comparison of three mixed dentition analysis methods in orthodontic patients at AKUH. J Coll Physicians Surg Pak. 2010 Aug; 20(8): 533-37.
- 60 Tome W, Ohyama Y, Yagi M, Takada K. Demonstration of a sex difference in the predictability of widths of unerupted permanent canines and premolars in a Japanese population. Angle Orthod. 2011 Nov; 81(6): 938-44.
- 61 van der Merwe SW, Rossouw P, van Wyk Kotze TJ, Trutero H. An adaptation of the Moyers mixed dentition space analysis for a Western Cape Caucasian population. J Dent Assoc S Afr. 1991 Sep; 46(9): 475-79.
- 62 Yuen KK, Tang EL, So LL. Mixed dentition analysis for Hong Kong Chinese. Angle Orthod. 1998 Feb; 68(1): 21-28.
- 63 Legovic M, Novosel A, Legovic A. Regression equations for determining mesiodistal crown diameters of canines and premolars. Angle Orthod. 2003 Jun; 73(3): 314-18.
- 64 Malkoc S, Basciftei FA, Nur M, Catalbas B. Maxillary and mandibular mesiodistal tooth sizes among different malocclusions in a sample of the Turkish population. Eur J Orthod. 2011 Oct; 33(5): 592-96.
- 65 Foneseca C. Predicting of mesiodistal crown width of the canine premolars segment in maxillary dental arches. Memphis, Tenn: University of Tennessee School of Dentistry; 1961.
- 66 Nourallah AW, Gesch D, Khordaji MN, Splieth C. New regression equations for predicting the size of unerupted canines and premolars in a contemporary population. Angle Orthod. 2002 Jun; 72(3): 216-21.
- 67 Melgaco CA, de Sousa Araujo MT, de Oliveira Ruellas AC. Mandibular permanent first molar and incisor width as predictor of mandibular canine and premolar width. Am J Orthod Dentofacial Orthop. 2007 Sep; 132(3): 340-45.
- 68 Cattaneo C, Butti AC, Bernini S, Biagi R, Salvato A. Comparative evaluation of the group of teeth with the best prediction value in the mixed dentition analysis. Eur J Paediatr Dent. 2010 Mar; 11(1): 23-26.
- 69 Ghulam Rasool Ub IUK, Noeen Arshad,, Owais Khalid Durrani, sohrab Shaheed. Applicability of Melgaco Equations for Predicting The Size of Unerupted Mandibular Canines and Premolars in Patients Reporting to Islamic International dental Hospital, Islamabad. Pakistan Oral & Dental Journal. 2008; 28(2): 165-70.